Can we calculate things better?

Can we calculate things better?

What does our solver look like now?

$$\boldsymbol{x}(t+\Delta t) = \boldsymbol{x}(t) + \dot{\boldsymbol{x}}\,\Delta t \quad + O(\Delta t^2)$$
$$\dot{\boldsymbol{x}}(t+\Delta t) = \dot{\boldsymbol{x}}(t) + \boldsymbol{a}(t)\,\Delta t \quad + O(\Delta t^2)$$

We say “first order solver” since errors are on order of Δt^2

Can we calculate things better?

What does our solver look like now?

$$\boldsymbol{x}(t+\Delta t) = \boldsymbol{x}(t) + \dot{\boldsymbol{x}}\,\Delta t \quad + O(\Delta t^2)$$
$$\dot{\boldsymbol{x}}(t+\Delta t) = \dot{\boldsymbol{x}}(t) + \boldsymbol{a}(t)\,\Delta t \quad + O(\Delta t^2)$$

We say “first order solver” since errors are on order of Δt^2

Hermite Solver

Can we calculate things better?

What does our solver look like now?

$$\boldsymbol{x}(t+\Delta t) = \boldsymbol{x}(t) + \dot{\boldsymbol{x}}\,\Delta t \quad + O(\Delta t^2)$$
$$\dot{\boldsymbol{x}}(t+\Delta t) = \dot{\boldsymbol{x}}(t) + \boldsymbol{a}(t)\,\Delta t \quad + O(\Delta t^2)$$

We say “first order solver” since errors are on order of Δt^2

Hermite Solver

$$\boldsymbol{x}_1 = \boldsymbol{x}_0 + \tfrac{1}{2}(\dot{\boldsymbol{x}}_1 + \dot{\boldsymbol{x}}_0)\Delta t + \tfrac{1}{12}(\boldsymbol{a}_0 - \boldsymbol{a}_1)\Delta t^2 \; + \mathcal{O}(\Delta t^5)$$
$$\dot{\boldsymbol{x}}_1 = \dot{\boldsymbol{x}}_0 + \tfrac{1}{2}(\boldsymbol{a}_1 + \boldsymbol{a}_0)\Delta t + \tfrac{1}{12}(\dot{\boldsymbol{a}}_0 - \dot{\boldsymbol{a}}_1)\Delta t^2 \; + \mathcal{O}(\Delta t^5)$$

4th order solver

Can we calculate things better?

What does our solver look like now?

$$\boldsymbol{x}(t+\Delta t) = \boldsymbol{x}(t) + \dot{\boldsymbol{x}}\,\Delta t \quad + O(\Delta t^2)$$
$$\dot{\boldsymbol{x}}(t+\Delta t) = \dot{\boldsymbol{x}}(t) + \boldsymbol{a}(t)\,\Delta t \quad + O(\Delta t^2)$$

We say “first order solver” since errors are on order of Δt^2

Hermite Solver

$$\boldsymbol{x}_1 = \boldsymbol{x}_0 + \tfrac{1}{2}(\dot{\boldsymbol{x}}_1 + \dot{\boldsymbol{x}}_0)\Delta t + \tfrac{1}{12}(\boldsymbol{a}_0 - \boldsymbol{a}_1)\Delta t^2 + \mathcal{O}(\Delta t^5)$$
$$\dot{\boldsymbol{x}}_1 = \dot{\boldsymbol{x}}_0 + \tfrac{1}{2}(\boldsymbol{a}_1 + \boldsymbol{a}_0)\Delta t + \tfrac{1}{12}(\dot{\boldsymbol{a}}_0 - \dot{\boldsymbol{a}}_1)\Delta t^2 + \mathcal{O}(\Delta t^5)$$

4th order solver

$$\boldsymbol{x}_p = \boldsymbol{x}_0 + \dot{\boldsymbol{x}}_0\Delta t + \tfrac{1}{2}\boldsymbol{a}_0\Delta t^2 + \tfrac{1}{6}\dot{\boldsymbol{a}}_0\Delta t^3$$
$$\dot{\boldsymbol{x}}_p = \dot{\boldsymbol{x}}_0 + \boldsymbol{a}_0\Delta t + \tfrac{1}{2}\dot{\boldsymbol{a}}_0\Delta t^2$$

Need “predictive” step for a and “jerk” $\dot{a}$ otherwise this is circular (need x_1 for a_1, but need a_1 for x_1…)

Can we calculate things better?

What does our solver look like now?

$$\boldsymbol{x}(t+\Delta t) = \boldsymbol{x}(t) + \dot{\boldsymbol{x}}\,\Delta t \quad + O(\Delta t^2)$$
$$\dot{\boldsymbol{x}}(t+\Delta t) = \dot{\boldsymbol{x}}(t) + \boldsymbol{a}(t)\,\Delta t \quad + O(\Delta t^2)$$

We say “first order solver” since errors are on order of Δt^2

Hermite Solver

$$\boldsymbol{x}_1 = \boldsymbol{x}_0 + \tfrac{1}{2}(\dot{\boldsymbol{x}}_1 + \dot{\boldsymbol{x}}_0)\Delta t + \tfrac{1}{12}(\boldsymbol{a}_0 - \boldsymbol{a}_1)\Delta t^2 + \mathcal{O}(\Delta t^5)$$
$$\dot{\boldsymbol{x}}_1 = \dot{\boldsymbol{x}}_0 + \tfrac{1}{2}(\boldsymbol{a}_1 + \boldsymbol{a}_0)\Delta t + \tfrac{1}{12}(\dot{\boldsymbol{a}}_0 - \dot{\boldsymbol{a}}_1)\Delta t^2 + \mathcal{O}(\Delta t^5)$$

4th order solver

$$\boldsymbol{x}_p = \boldsymbol{x}_0 + \dot{\boldsymbol{x}}_0\Delta t + \tfrac{1}{2}\boldsymbol{a}_0\Delta t^2 + \tfrac{1}{6}\dot{\boldsymbol{a}}_0\Delta t^3$$
$$\dot{\boldsymbol{x}}_p = \dot{\boldsymbol{x}}_0 + \boldsymbol{a}_0\Delta t + \tfrac{1}{2}\dot{\boldsymbol{a}}_0\Delta t^2$$

Need “predictive” step for a and “jerk” $\dot{a}$ otherwise this is circular (need x_1 for a_1, but need a_1 for x_1…)

www.astroblend.com/ba2016/extra_resources/integrators.pdf

Can we calculate things better?

One more thing: “Gravitational Units”

Basically: G = M = R = 1

N-body Units

A conventional system of units in which

$$\begin{aligned} G &= 1 \\ M &= 1 \\ R &= 1 \end{aligned}$$

Example

Suppose a star cluster has $M = 10^5 M_\odot, R = 5\text{pc}$. To convert a velocity from the N-body code to km/s, multiply by $\sqrt{\frac{GM}{R}}$, where G is expressed in the same units (i.e. km/s, $M_\odot$, pc), i.e. $G \simeq 0.043$.

(slides brazenly stolen from Heggie)

On day 2 website:
Some code that compares Hermite and Euler Integration Schemes with Analytical

www.astroblend.com/ba2016/code/day2/twobodyintegrators_eu_and_hermite_jills.py

Off to N-Body! Lets talk about smashing planets…

Super Planet Crash! http://www.stefanom.org/spc/

Bonus - dealing with alien overlords:
http://save-point.herokuapp.com/dashboard/users.php